

AN EXISTENCE THEOREM FOR THE YAMABE PROBLEM ON MANIFOLDS WITH BOUNDARY

SIMON BRENDLE AND SZU-YU SOPHIE CHEN

1. INTRODUCTION

The Yamabe problem, solved by Trudinger [14], Aubin [1], and Schoen [12], asserts that any Riemannian metric on a closed manifold is conformal to a metric with constant scalar curvature. Escobar [8], [9] has studied analogous questions on manifolds with boundary. To fix notation, let (M, g) be a compact Riemannian manifold of dimension $n \geq 3$ with boundary ∂M . We denote by R_g the scalar curvature of (M, g) and by κ_g the mean curvature of the boundary ∂M . There are two natural ways to extend the Yamabe problem to manifolds with boundary:

- (a) Find a metric $\tilde{g}$ in the conformal class of g such that $R_{\tilde{g}}$ is constant and $\kappa_{\tilde{g}} = 0$.
- (b) Find a metric $\tilde{g}$ in the conformal class of g such that $R_{\tilde{g}} = 0$ and $\kappa_{\tilde{g}}$ is constant.

The boundary value problem (a) was first proposed by Escobar [8]. The boundary value problem (b) is studied in [9] and [11].

In this paper, we focus on the boundary value problem (a). The solvability of (a) is equivalent to the existence of a critical point of the Yamabe functional. This functional is defined by

$$E_g(u) = \frac{\int_M \left(\frac{4(n-1)}{n-2} |du|_g^2 + R_g u^2 \right) d\text{vol}_g + \int_{\partial M} 2\kappa_g u^2 d\sigma_g}{\left(\int_M u^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}}},$$

where u is a smooth positive function on M . Moreover, the Yamabe constant is defined as

$$Y(M, \partial M, g) = \inf_{0 < u \in C^\infty(M)} E_g(u).$$

It is well known that $Y(M, \partial M, g)$ is invariant under a conformal change of the metric g . Moreover, $Y(M, \partial M, g) \leq Y(S_+^n, \partial S_+^n)$, where $Y(S_+^n, \partial S_+^n)$ denotes the Yamabe constant of the hemisphere S_+^n equipped with the standard metric.

The first author was partially supported by a Brown foundation fellowship and by the National Science Foundation under grant DMS-0905628. The second author was supported in part by the Miller Institute for Basic Research in Science and the National Science Foundation.

Proving the existence of a minimizer for the functional E_g is a difficult problem, as E_g does not satisfy the Palais-Smale condition. The following existence result was established by Escobar [8].

Theorem 1 (J. Escobar [8]). *If $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$, then there exists a metric $\tilde{g}$ in the conformal class of g such that $R_{\tilde{g}}$ is constant and $\kappa_{\tilde{g}}$ is equal to 0.*

Theorem 1 should be compared to Aubin's existence theorem for the Yamabe problem on manifolds without boundary (cf. [1]).

In dimension $3 \leq n \leq 5$, Escobar showed that $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$ unless M is conformally equivalent to the hemisphere S_+^n . In dimension $n \geq 6$, Escobar was able to verify the inequality $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$ under the assumption that the boundary ∂M is not umbilic.

Therefore, it remains to consider the case that $n \geq 6$ and ∂M is umbilic. For abbreviation, we put $d = \lfloor \frac{n-2}{2} \rfloor$. As in [4], we denote by $\mathcal{Z}$ the set of all points $p \in M$ such that

$$\limsup_{x \rightarrow p} d(p, x)^{2-d} |W_g(x)| = 0,$$

where W_g denotes the Weyl tensor of (M, g) . In other words, a point $p \in M$ belongs to $\mathcal{Z}$ if and only if $D^m W_g(p) = 0$ for all $m \in \{0, 1, \dots, d-2\}$. Note that the set $\mathcal{Z}$ is invariant under a conformal change of the metric.

The following is the main result of this paper:

Theorem 2. *Let (M, g) be a compact Riemannian manifold of dimension $n \geq 6$ with umbilic boundary ∂M . Moreover, let $p \in \partial M$ be an arbitrary point on the boundary of M . If $p \notin \mathcal{Z}$, then $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$. Consequently, there exists a metric $\tilde{g}$ in the conformal class of g such that $R_{\tilde{g}}$ is constant and $\kappa_{\tilde{g}}$ is equal to 0.*

In case $p \in \mathcal{Z}$, we are able to show that $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$, provided that a certain asymptotically flat manifold has positive ADM mass (see Theorem 20 below).

We now give an outline of the proof of Theorem 2. By a theorem of Marques [11], we may work in conformal Fermi coordinates around p . We define a function u_ε by

$$(1) \quad u_\varepsilon(x) = \left(\frac{\varepsilon}{\varepsilon^2 + |x|^2} \right)^{\frac{n-2}{2}}.$$

The function u_ε satisfies

$$\Delta u_\varepsilon = -n(n-2) u_\varepsilon^{\frac{n+2}{n-2}}$$

and

$$u_\varepsilon \partial_i \partial_k u_\varepsilon - \frac{n}{n-2} \partial_i u_\varepsilon \partial_k u_\varepsilon = \frac{1}{n} \left(u_\varepsilon \Delta u_\varepsilon - \frac{n}{n-2} |du_\varepsilon|^2 \right) \delta_{ik}.$$

These identities reflect the fact that the metric $u_\varepsilon^{\frac{4}{n-2}} \delta_{ik}$ is Einstein.

We then consider a sum of the form $u_\varepsilon + w$, where u_ε is given by (1) and w is a correction term. This function is only defined in a small neighborhood of the point p . In order to extend the testfunction to all of M , we glue the function $u_\varepsilon + w$ to the Greens function of the conformal Laplacian with pole at p .

In order to show that the resuling testfunction has Yamabe energy less than $Y(S_+^n, \partial S_+^n)$, we make extensive use of techniques developed in [4] (see also [5], [6], [7]). In [4], these techniques were used to prove a convergence theorem for the parabolic Yamabe flow in dimension $n \geq 6$. The convergence of the Yamabe flow in dimension $3 \leq n \leq 5$ was shown in [3].

2. AUXILIARY RESULTS

In this section, we consider the halfspace $\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n \geq 0\}$. Moreover, we assume that $H_{ik}(x)$ is a trace-free symmetric two-tensor on $\mathbb{R}_+^n$ which satisfies the following conditions:

- At each point $x \in \mathbb{R}_+^n$, we have $H_{in}(x) = 0$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\sum_{k=1}^n H_{ik}(x) x_k = 0$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\partial_n H_{ik}(x) = 0$ for all $i, k \in \{1, \dots, n\}$.

Finally, we assume that the components $H_{ik}(x)$ are polynomials of the form

$$H_{ik}(x) = \sum_{2 \leq |\alpha| \leq d} h_{ik,\alpha} x^\alpha,$$

where the sum is taken over all multi-indices α of length $2 \leq |\alpha| \leq d$.

As in [4], we define

$$A_{ik} = \sum_{m=1}^n \partial_i \partial_m H_{mk} + \sum_{m=1}^n \partial_m \partial_k H_{im} - \Delta H_{ik} - \frac{1}{n-1} \sum_{m,p=1}^n \partial_m \partial_p H_{mp} \delta_{ik}$$

and

$$\begin{aligned} Z_{ijkl} &= \partial_i \partial_k H_{jl} - \partial_i \partial_l H_{jk} - \partial_j \partial_k H_{il} + \partial_j \partial_l H_{ik} \\ &\quad + \frac{1}{n-2} (A_{jl} \delta_{ik} - A_{jk} \delta_{il} - A_{il} \delta_{jk} + A_{ik} \delta_{jl}). \end{aligned}$$

Note that

$$\partial_l Z_{ijkl} = \frac{n-3}{n-2} (\partial_i A_{jk} - \partial_j A_{ik}).$$

Lemma 3. *We have $A_{in}(x) = 0$ for all points $x \in \partial\mathbb{R}_+^n$ and all indices $i \in \{1, \dots, n-1\}$.*

Proof. Note that $\partial_n H_{im}(x) = 0$ for all points $x \in \partial\mathbb{R}_+^n$ and all $i, m \in \{1, \dots, n-1\}$. If we differentiate this identity in tangential direction, we obtain

$$A_{in}(x) = \sum_{m=1}^{n-1} \partial_m \partial_n H_{im}(x) = 0$$

for all points $x \in \partial\mathbb{R}_+^n$ and all $i \in \{1, \dots, n-1\}$.

Lemma 4. *Assume that $Z_{ijkl}(x) = 0$ for all points $x \in \partial\mathbb{R}_+^n$ and all $i, j, k, l \in \{1, \dots, n\}$. Then $H_{ik}(x) = A_{ik}(x) = 0$ for all $x \in \partial\mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$.*

Proof. We define

$$\begin{aligned} \hat{A}_{ik}(x) &= \sum_{m=1}^{n-1} \partial_i \partial_m H_{mk}(x) + \sum_{m=1}^{n-1} \partial_m \partial_k H_{im}(x) \\ &\quad - \sum_{m=1}^{n-1} \partial_m \partial_m H_{ik}(x) - \frac{1}{n-2} \sum_{m,p=1}^n \partial_m \partial_p H_{mp}(x) \delta_{ik} \end{aligned}$$

for all points $x \in \partial\mathbb{R}_+^n$ and all indices $i, k \in \{1, \dots, n-1\}$. By assumption, we have

$$\partial_n \partial_n H_{ik}(x) + \frac{1}{n-2} (A_{ik}(x) + A_{nn}(x) \delta_{ik}) = Z_{inkn}(x) = 0$$

for all points $x \in \partial\mathbb{R}_+^n$ and all indices $i, k \in \{1, \dots, n-1\}$. This implies

$$\begin{aligned} \hat{A}_{ik}(x) &= A_{ik}(x) + \partial_n \partial_n H_{ik}(x) - \frac{1}{(n-1)(n-2)} \sum_{m,p=1}^{n-1} \partial_m \partial_p H_{mp}(x) \delta_{ik} \\ &= A_{ik}(x) + \partial_n \partial_n H_{ik}(x) + \frac{1}{n-2} A_{nn}(x) \delta_{ik} \\ &= \frac{n-3}{n-2} A_{ik}(x) \end{aligned}$$

for all points $x \in \partial\mathbb{R}_+^n$ and all indices $i, k \in \{1, \dots, n-1\}$. Hence, we obtain

$$\begin{aligned} &\partial_i \partial_k H_{jl}(x) - \partial_i \partial_l H_{jk}(x) - \partial_j \partial_k H_{il}(x) + \partial_j \partial_l H_{ik}(x) \\ &= -\frac{1}{n-2} (A_{jl}(x) \delta_{ik} - A_{jk}(x) \delta_{il} - A_{il}(x) \delta_{jk} + A_{ik}(x) \delta_{jl}) \\ &= -\frac{1}{n-3} (\hat{A}_{jl}(x) \delta_{ik} - \hat{A}_{jk}(x) \delta_{il} - \hat{A}_{il}(x) \delta_{jk} + \hat{A}_{ik}(x) \delta_{jl}) \end{aligned}$$

for all points $x \in \partial\mathbb{R}_+^n$ and all indices $i, j, k, l \in \{1, \dots, n-1\}$. Using Proposition 7 in [4], we conclude that $H_{ik}(x) = 0$ for all $x \in \partial\mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$. This implies $A_{ik}(x) = \frac{n-2}{n-3} \hat{A}_{ik}(x) = 0$ for all $x \in \partial\mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$.

Proposition 5. *Assume that $Z_{ijkl}(x) = 0$ for all $x \in \mathbb{R}_+^n$ and all $i, j, k, l \in \{1, \dots, n\}$. Then $H_{ik}(x) = 0$ for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$.*

Proof. Without loss of generality, we may assume that $H_{ik}(x)$ is homogenous of degree $d' \geq 2$. By assumption, we have $Z_{ijkl}(x) = 0$ for all

$x \in \mathbb{R}_+^n$ and all $i, j, k, l \in \{1, \dots, n\}$. This implies

$$\partial_i A_{jk}(x) - \partial_j A_{ik}(x) = \frac{n-2}{n-3} \sum_{l=1}^n \partial_l Z_{ijkl}(x) = 0$$

for all $x \in \mathbb{R}_+^n$ and all $i, j, k \in \{1, \dots, n\}$. We next define

$$\varphi(x) = \frac{1}{d'(d'-1)} \sum_{i,k=1}^n A_{ik}(x) x_i x_k$$

for all $x \in \mathbb{R}_+^n$. Clearly, $\varphi(x)$ is a homogeneous polynomial of degree d' . Moreover, we have

$$\partial_k \varphi(x) = \frac{1}{d'-1} \sum_{i=1}^n A_{ik}(x) x_i$$

for all $x \in \mathbb{R}_+^n$ and all $k \in \{1, \dots, n\}$. This implies

$$\partial_i \partial_k \varphi(x) = A_{ik}(x)$$

for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n\}$. Using the identity $\sum_{i=1}^{n-1} H_{ii}(x) = 0$, we obtain

$$\partial_n \partial_n \varphi(x) = A_{nn}(x) = -\frac{1}{n-1} \sum_{i=1}^{n-1} A_{ii}(x) = -\frac{1}{n-1} \sum_{i=1}^{n-1} \partial_i \partial_i \varphi(x)$$

for all $x \in \mathbb{R}_+^n$. By Lemma 3, we have $\partial_n \varphi(x) = \frac{1}{d'-1} \sum_{i=1}^{n-1} A_{in}(x) x_i = 0$ for all $x \in \partial \mathbb{R}_+^n$. Moreover, it follows from Lemma 4 that $\varphi(x) = \frac{1}{d'(d'-1)} \sum_{i,k=1}^{n-1} A_{ik}(x) x_i x_k = 0$ for all $x \in \partial \mathbb{R}_+^n$. Putting these facts together, we conclude that $\varphi(x) = 0$ for all $x \in \mathbb{R}_+^n$.

Therefore, we have $A_{ik}(x) = \partial_i \partial_k \varphi(x) = 0$ for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n\}$. This implies

$$\partial_n \partial_n H_{ik}(x) = Z_{inkn}(x) - \frac{1}{n-2} (A_{ik}(x) + A_{nn}(x) \delta_{ik}) = 0$$

for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$. On the other hand, we know that $H_{ik}(x) = \partial_n H_{ik}(x) = 0$ for all $x \in \partial \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$. From this, it follows that $H_{ik}(x) = 0$ for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$. This completes the proof of Proposition 5.

For each $r > 0$, we denote by $U_r \subset \mathbb{R}_+^n$ the open ball of radius $\frac{r}{4}$ centered at the point $(0, \dots, 0, \frac{3r}{2})$. Moreover, let $u_\varepsilon : \mathbb{R}_+^n \rightarrow \mathbb{R}$ be defined by (1).

Proposition 6. *There exists a constant K_1 , depending only on n , such that*

$$\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 r^{2|\alpha|-4+n} \leq K_1 \int_{U_r} \sum_{i,j,k,l=1}^n |Z_{ijkl}(x)|^2 dx$$

for all $r > 0$.

Proof. It follows from Proposition 5 that the assertion holds for $r = 1$. The general case follows by scaling.

Proposition 7. *Let V be a smooth vector field on $\mathbb{R}_+^n$. Moreover, let*

$$T_{ik} = H_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}$$

and

$$\begin{aligned} Q_{ik,l} &= u_\varepsilon \partial_l T_{ik} - \frac{2}{n-2} \partial_i u_\varepsilon T_{kl} - \frac{2}{n-2} \partial_k u_\varepsilon T_{il} \\ &\quad + \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{ip} \delta_{kl} + \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{kp} \delta_{il}. \end{aligned}$$

Then there exists a constant K_2 , depending only on n , such that

$$\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} r^{2|\alpha|+2-n} \leq K_2 \int_{(B_{2r}(0) \setminus B_r(0)) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n |Q_{ik,l}(x)|^2 dx$$

for all $r \geq \varepsilon$.

Proof. In [4], the first author showed that

$$\begin{aligned} \frac{1}{4} \sum_{i,j,k,l=1}^n |Z_{ijkl}|^2 &= \sum_{i,j,k,l=1}^n \partial_j (u_\varepsilon^{-1} Q_{ik,l}) Z_{ijkl} \\ &\quad + \frac{2}{n-2} \sum_{i,j,k,l=1}^n u_\varepsilon^{-2} \partial_k u_\varepsilon Q_{il,j} Z_{ijkl} \end{aligned}$$

(cf. [4], p. 555). Let us fix a smooth cut-off function $\eta : \mathbb{R}^n \rightarrow [0, 1]$ such that $\eta(x) = 1$ for $x \in U_1$ and $\eta(x) = 0$ for $x \notin (B_2(0) \setminus B_1(0)) \cap \mathbb{R}_+^n$. In particular, we have $\eta(x) = 0$ for all $x \in \partial \mathbb{R}_+^n$. Integration by parts gives

$$\begin{aligned} &\int_{\mathbb{R}_+^n} \frac{1}{4} \sum_{i,j,k,l=1}^n |Z_{ijkl}(x)|^2 \eta(x/r) dx \\ &= - \int_{\mathbb{R}_+^n} \sum_{i,j,k,l=1}^n u_\varepsilon(x)^{-1} Q_{ik,l}(x) \partial_j [Z_{ijkl}(x) \eta(x/r)] dx \\ &\quad + \int_{\mathbb{R}_+^n} \frac{2}{n-2} \sum_{i,j,k,l=1}^n u_\varepsilon(x)^{-2} \partial_k u_\varepsilon(x) Q_{il,j}(x) Z_{ijkl}(x) \eta(x/r) dx. \end{aligned}$$

Using Hölder's inequality, we obtain

$$\begin{aligned} & \int_{U_r} \sum_{i,j,k,l=1}^n |Z_{ijkl}(x)|^2 dx \\ & \leq K_3 \varepsilon^{-\frac{n-2}{2}} r^{n-3} \left(\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 r^{2|\alpha|-4+n} \right)^{\frac{1}{2}} \\ & \quad \cdot \left(\int_{(B_{2r} \setminus B_r(0)) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n |Q_{ik,l}(x)|^2 dx \right)^{\frac{1}{2}} \end{aligned}$$

for all $r \geq \varepsilon$. Here, K_3 is a positive constant that depends only on n . On the other hand, it follows from Proposition 6 that

$$\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 r^{2|\alpha|-4+n} \leq K_1 \int_{U_r} \sum_{i,j,k,l=1}^n |Z_{ijkl}(x)|^2 dx.$$

Putting these facts together, the assertion follows.

Corollary 8. *Let V be a smooth vector field on $\mathbb{R}_+^n$. Moreover, let*

$$T_{ik} = H_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}$$

and

$$\begin{aligned} Q_{ik,l} &= u_\varepsilon \partial_l T_{ik} - \frac{2}{n-2} \partial_i u_\varepsilon T_{kl} - \frac{2}{n-2} \partial_k u_\varepsilon T_{il} \\ &+ \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{ip} \delta_{kl} + \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{kp} \delta_{il}. \end{aligned}$$

Then there exists a constant K_4 , depending only on n , such that

$$\begin{aligned} & \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\ & \leq K_4 \int_{B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n |Q_{ik,l}(x)|^2 dx \end{aligned}$$

for all $\delta \geq 2\varepsilon$.

3. THE MAIN ESTIMATE

We now describe the construction of the test function. Let (M, g) be a compact Riemannian manifold of dimension $n \geq 6$ with umbilic boundary ∂M . After changing the metric conformally, we may assume that ∂M is totally geodesic.

Let us fix a point $p \in \partial M$, and let $(x_1, \dots, x_n)$ denote the Fermi coordinates around p . In these coordinates, the metric has the following properties:

- At each point $x \in \mathbb{R}_+^n$, we have $g_{in}(x) = \delta_{in}$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\sum_{k=1}^n g_{ik}(x) x_k = x_i$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\partial_n g_{ik}(x) = 0$ for all $i, k \in \{1, \dots, n\}$.

By a theorem of Marques, there exists a system of conformal Fermi coordinates around p (see [11], Proposition 3.1). Hence, after performing a conformal change of the metric, we may assume that $\det g(x) = 1 + O(|x|^{2d+2})$, where $d = \lfloor \frac{n-2}{2} \rfloor$.

In the next step, we write $g(x) = \exp(h(x))$, where $h(x)$ is a smooth function taking values in the space of symmetric $n \times n$ matrices. This function has the following properties:

- At each point $x \in \mathbb{R}_+^n$, we have $h_{in}(x) = 0$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\sum_{k=1}^n h_{ik}(x) x_k = 0$ for all $i \in \{1, \dots, n\}$.
- At each point $x \in \partial\mathbb{R}_+^n$, we have $\partial_n h_{ik}(x) = 0$ for all $i, k \in \{1, \dots, n\}$.

Moreover, we have $\text{tr } h(x) = O(|x|^{2d+2})$. For abbreviation, we denote by

$$H_{ik}(x) = \sum_{2 \leq |\alpha| \leq d} h_{ik,\alpha} x^\alpha$$

the Taylor polynomial of order d associated with the function $h_{ik}(x)$. Clearly, $H_{ik}(x)$ is a trace-free symmetric two-tensor on $\mathbb{R}_+^n$. Moreover, we have $h_{ik}(x) = H_{ik}(x) + O(|x|^{d+1})$.

Let us fix a non-negative smooth function such that $\chi(t) = 1$ for $t \leq \frac{4}{3}$ and $\chi(t) = 0$ for $t \geq \frac{5}{3}$. Given any $\delta > 0$, we define a cut-off function $\chi_\delta : \mathbb{R}^n \rightarrow \mathbb{R}$ by $\chi_\delta(x) = \chi(|x|/\delta)$. By Theorem 24, there exists a smooth vector field V on $\mathbb{R}_+^n$ with the following properties:

- At each point $x \in \mathbb{R}_+^n$, we have

$$\sum_{k=1}^n \partial_k \left[u_\varepsilon^{\frac{2n}{n-2}} (\chi_\delta H_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \text{div } V \delta_{ik}) \right] = 0$$

for all $i \in \{1, \dots, n\}$.

- At each point $x \in \partial\mathbb{R}_+^n$, we have $V_n(x) = \partial_n V_i(x) = 0$ for all $i \in \{1, \dots, n-1\}$.

By Corollary 26, the vector field V satisfies the estimate

$$(2) \quad |\partial^\beta V^{(\varepsilon, \delta)}(x)| \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| (\varepsilon + |x|)^{|\alpha|+1-|\beta|}$$

for every multi-index β and all $x \in \mathbb{R}_+^n$. Here, C is a positive constant that depends only on n and $|\beta|$.

For abbreviation, we define

$$\begin{aligned} S_{ik} &= \partial_i V_k + \partial_k V_i - \frac{2}{n} \text{div } V \delta_{ik}, \\ T_{ik} &= H_{ik} - S_{ik}, \end{aligned}$$

$$\begin{aligned}
Q_{ik,l} &= u_\varepsilon \partial_l T_{ik} - \frac{2}{n-2} \partial_i u_\varepsilon T_{kl} - \frac{2}{n-2} \partial_k u_\varepsilon T_{il} \\
&\quad + \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{ip} \delta_{kl} + \frac{2}{n-2} \sum_{p=1}^n \partial_p u_\varepsilon T_{kp} \delta_{il},
\end{aligned}$$

and

$$w = \sum_{l=1}^n \partial_l u_\varepsilon V_l + \frac{n-2}{2n} u_\varepsilon \operatorname{div} V.$$

By definition of V , we have

$$(3) \quad \sum_{k=1}^n \partial_k (u_\varepsilon^{\frac{2n}{n-2}} T_{ik}) = 0$$

for all points $x \in B_\delta(0) \cap \mathbb{R}_+^n$ and all $i \in \{1, \dots, n\}$. This implies

$$(4) \quad \sum_{k=1}^n \left(u_\varepsilon \partial_k T_{ik} + \frac{2n}{n-2} \partial_k u_\varepsilon T_{ik} \right) = 0$$

for all points $x \in B_\delta(0) \cap \mathbb{R}_+^n$ and all $i \in \{1, \dots, n\}$. The following result was established in [4]:

Proposition 9 (S. Brendle [4]). *There exists a smooth vector field ξ on $\mathbb{R}_+^n$ such that*

$$\begin{aligned}
&\frac{1}{4} u_\varepsilon^2 \sum_{i,k,l=1}^n \partial_l H_{ik} \partial_l H_{ik} - \frac{1}{2} u_\varepsilon^2 \sum_{i,k,l=1}^n \partial_k H_{ik} \partial_l H_{il} \\
&- 2 u_\varepsilon \sum_{i,k,l=1}^n \partial_k u_\varepsilon H_{ik} \partial_l H_{il} - \frac{2(n-1)}{n-2} \sum_{i,k,l=1}^n \partial_k u_\varepsilon \partial_l u_\varepsilon H_{ik} H_{il} \\
&- 2 u_\varepsilon w \sum_{i,k=1}^n \partial_i \partial_k H_{ik} + \frac{8(n-1)}{n-2} \sum_{i,k=1}^n \partial_i u_\varepsilon \partial_k w H_{ik} \\
&- \frac{4(n-1)}{n-2} |dw|^2 + \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
&= \frac{1}{4} \sum_{i,k,l=1}^n Q_{ik,l} Q_{ik,l} + 2 u_\varepsilon^{\frac{2n}{n-2}} \sum_{i,k=1}^n T_{ik} T_{ik} + \operatorname{div} \xi
\end{aligned}$$

for all points $x \in B_\delta(0) \cap \mathbb{R}_+^n$.

The vector field ξ can be expressed in terms of the tensor H_{ik} and the vector field V (cf. [4], Section 2). In the next step, we show that ξ is tangential along $\partial \mathbb{R}_+^n$. To that end, we need the following lemma:

Lemma 10. *At each point $x \in B_\delta(0) \cap \partial \mathbb{R}_+^n$, we have*

$$S_{in}(x) = T_{in}(x) = 0$$

and

$$\partial_n S_{ik}(x) = \partial_n T_{ik}(x) = 0$$

for all $i, k \in \{1, \dots, n-1\}$. Moreover, we have $\partial_n S_{nn}(x) = \partial_n T_{nn}(x) = 0$ and $\partial_n w(x) = 0$.

Proof. By assumption, we have $V_n(x) = \partial_n V_i(x) = 0$ for all $x \in \partial \mathbb{R}_+^n$. This implies $S_{in}(x) = T_{in}(x) = 0$ for all $i \in \{1, \dots, n-1\}$. This implies

$$\sum_{k=1}^{n-1} \left(u_\varepsilon(x) \partial_k T_{kn}(x) + \frac{2n}{n-2} \partial_k u_\varepsilon(x) T_{kn}(x) \right) = 0.$$

Using (4), we obtain

$$u_\varepsilon(x) \partial_n T_{nn}(x) + \frac{2n}{n-2} \partial_n u_\varepsilon(x) T_{nn}(x) = 0.$$

This implies $\partial_n T_{nn}(x) = 0$, hence $\partial_n S_{nn}(x) = 0$. Consequently, we have $\partial_n \partial_n V(x) = 0$. From this, the assertion follows easily.

Lemma 11. *We have $\xi_n(x) = 0$ for all points $x \in B_\delta(0) \cap \partial \mathbb{R}_+^n$.*

Proof. The vector field ξ satisfies

$$\begin{aligned} \xi_n = & -2 \sum_{k=1}^n u_\varepsilon w \partial_k H_{nk} + 2 \sum_{i=1}^n \partial_i (u_\varepsilon w) H_{in} \\ & + \frac{1}{2} u_\varepsilon^2 \sum_{i,k=1}^n \partial_n S_{ik} H_{ik} - u_\varepsilon^2 \sum_{i,l=1}^n \partial_l S_{il} H_{in} - 2 u_\varepsilon \sum_{i,l=1}^n \partial_l u_\varepsilon S_{il} H_{in} \\ & + u_\varepsilon w \sum_{k=1}^n \partial_k S_{nk} - \sum_{i=1}^n \partial_i (u_\varepsilon w) S_{in} \\ & - \frac{1}{4} u_\varepsilon^2 \sum_{i,k=1}^n \partial_n S_{ik} S_{ik} + \frac{1}{2} u_\varepsilon^2 \sum_{i,l=1}^n \partial_l S_{il} S_{in} + u_\varepsilon \sum_{i,l=1}^n \partial_l u_\varepsilon S_{il} S_{in} \\ & + \frac{4(n-1)}{n-2} \sum_{i=1}^n \partial_i u_\varepsilon w S_{in} - \frac{4(n-1)}{n-2} w \partial_n w \\ & + \frac{2}{n-2} u_\varepsilon \sum_{i,k=1}^n \partial_k u_\varepsilon T_{ik} T_{in} \end{aligned}$$

(see [4], Section 2). Using Lemma 10, we conclude that $\xi_n(x) = 0$ for all $x \in B_\delta(0) \cap \partial \mathbb{R}_+^n$.

Proposition 12. *We have*

$$\begin{aligned}
& \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left[\frac{1}{4} u_\varepsilon^2 \sum_{i,k,l=1}^n \partial_l H_{ik} \partial_l H_{ik} - \frac{1}{2} u_\varepsilon^2 \sum_{i,k,l=1}^n \partial_k H_{ik} \partial_l H_{il} \right] \\
& - \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left[2 u_\varepsilon \sum_{i,k,l=1}^n \partial_k u_\varepsilon H_{ik} \partial_l H_{il} + \frac{2(n-1)}{n-2} \sum_{i,k,l=1}^n \partial_k u_\varepsilon \partial_l u_\varepsilon H_{ik} H_{il} \right] \\
& - \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left[2 u_\varepsilon w \sum_{i,k=1}^n \partial_i \partial_k H_{ik} - \frac{8(n-1)}{n-2} \sum_{i,k=1}^n \partial_i u_\varepsilon \partial_k w H_{ik} \right] \\
& - \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left[\frac{4(n-1)}{n-2} |dw|^2 - \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \right] \\
& \geq 2\lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& - C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{n-2}.
\end{aligned}$$

for $\delta \geq 2\varepsilon$. Here, λ and C are positive constants that depend only on n .

Proof. We consider the identity in Proposition 9 and integrate over $B_\delta(0) \cap \mathbb{R}_+^n$. By Corollary 8, we have

$$\begin{aligned}
& \int_{B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n Q_{ik,l} Q_{ik,l} \\
& \geq 8\lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+3-2n} dx,
\end{aligned}$$

where $\lambda = 1/(8K_4)$ is a positive constant that depends only on n . Moreover, it follows from 2 that

$$|\xi(x)| \leq C \varepsilon^{n-2} \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| (\varepsilon + |x|)^{2|\alpha|+3-2n}$$

for all $x \in \mathbb{R}_+^n$. Using Lemma 11 and the divergence theorem, we obtain

$$\begin{aligned}
\int_{B_\delta(0) \cap \mathbb{R}_+^n} \operatorname{div} \xi &= \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} \xi_i - \int_{B_\delta(0) \cap \partial \mathbb{R}_+^n} \xi_n \\
&\leq C \varepsilon^{n-2} \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \delta^{2|\alpha|+2-n}.
\end{aligned}$$

Putting these facts together, the assertion follows.

Finally, we need the following estimate for the scalar curvature R_g .

Proposition 13. *The scalar curvature R_g satisfies the estimates*

$$(5) \quad \left| R_g - \sum_{i,k=1}^n \partial_i \partial_k H_{ik} \right| \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| |x|^{|\alpha|} + C |x|^{d-1}$$

and

$$(6) \quad \begin{aligned} & \left| R_g - \sum_{i,k=1}^n \partial_i \partial_k h_{ik} + \sum_{i,k,l=1}^n \partial_k (H_{ik} \partial_l H_{il}) \right. \\ & \quad \left. - \frac{1}{2} \sum_{i,k,l=1}^n \partial_k H_{ik} \partial_l H_{il} + \frac{1}{4} \sum_{i,k,l=1}^n \partial_l H_{ik} \partial_l H_{ik} \right| \\ & \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 |x|^{2|\alpha|} \\ & \quad + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| |x|^{|\alpha|+d-1} + C |x|^{2d} \end{aligned}$$

if $|x|$ is sufficiently small.

Proof. This follows easily from Proposition 25 in [5] (see also Corollary 12 in [4], where geodesic normal coordinates are considered).

Our goal is to estimate the Yamabe energy of $u_\varepsilon + w$. To that end, we proceed in several steps:

Proposition 14. *There exist positive constants λ , C , δ_0 such that*

$$\begin{aligned} & \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left(\frac{4(n-1)}{n-2} |d(u_\varepsilon + w)|_g^2 + R_g (u_\varepsilon + w)^2 \right) \\ & \leq \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} |du_\varepsilon|^2 + \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\ & \quad + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k=1}^n \frac{x_i}{|x|} (u_\varepsilon^2 \partial_k h_{ik} - 2 u_\varepsilon \partial_k u_\varepsilon h_{ik}) \\ & \quad - \lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\ & \quad + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2} \end{aligned}$$

if $0 < 2\varepsilon \leq \delta \leq \delta_0$. The constant λ depends only on n . The constants C , δ_0 depend on the background manifold (M, g) .

Proof. Let us write

$$\begin{aligned}
& \frac{4(n-1)}{n-2} |d(u_\varepsilon + w)|_g^2 + R_g(u_\varepsilon + w)^2 \\
&= \frac{4(n-1)}{n-2} |du_\varepsilon|^2 + \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
&+ \frac{8(n-1)}{n-2} J^{(1)} + J^{(2)} + J^{(3)} + J^{(4)} + J^{(5)} + J^{(6)} + J^{(7)},
\end{aligned}$$

where

$$\begin{aligned}
J^{(1)} &= \sum_{i=1}^n \partial_i u_\varepsilon \partial_i w \\
J^{(2)} &= -\frac{4(n-1)}{n-2} \sum_{i,k=1}^n \partial_i u_\varepsilon \partial_k u_\varepsilon h_{ik} + u_\varepsilon^2 \sum_{i,k=1}^n \partial_i \partial_k h_{ik} \\
J^{(3)} &= -u_\varepsilon^2 \sum_{i,k,l=1}^n \partial_k (H_{ik} \partial_l H_{il}) - 2u_\varepsilon \sum_{i,k,l=1}^n \partial_k u_\varepsilon H_{ik} \partial_l H_{il} \\
J^{(4)} &= -\frac{1}{4} \sum_{i,k,l=1}^n u_\varepsilon^2 \partial_l H_{ik} \partial_l H_{ik} + \frac{1}{2} \sum_{i,k,l=1}^n u_\varepsilon^2 \partial_k H_{ik} \partial_l H_{il} \\
&+ 2u_\varepsilon \sum_{i,k,l=1}^n \partial_k u_\varepsilon H_{ik} \partial_l H_{il} + \frac{2(n-1)}{n-2} \sum_{i,k,l=1}^n \partial_k u_\varepsilon \partial_l u_\varepsilon H_{ik} H_{il} \\
&+ 2u_\varepsilon w \sum_{i,k=1}^n \partial_i \partial_k H_{ik} - \frac{8(n-1)}{n-2} \sum_{i,k=1}^n \partial_i u_\varepsilon \partial_k w H_{ik} \\
&+ \frac{4(n-1)}{n-2} |dw|^2 - \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
J^{(5)} &= \frac{4(n-1)}{n-2} \sum_{i,k=1}^n \left[g^{ik} - \delta_{ik} + h_{ik} - \frac{1}{2} \sum_{l=1}^n H_{il} H_{kl} \right] \partial_i u_\varepsilon \partial_k u_\varepsilon \\
&+ \left[R_g - \sum_{i,k=1}^n \partial_i \partial_k h_{ik} + \sum_{i,k,l=1}^n \partial_k (H_{ik} \partial_l H_{il}) \right. \\
&\quad \left. - \frac{1}{2} \sum_{i,k,l=1}^n \partial_k H_{ik} \partial_l H_{il} + \frac{1}{4} \sum_{i,k,l=1}^n \partial_l H_{ik} \partial_l H_{ik} \right] u_\varepsilon^2
\end{aligned}$$

and

$$\begin{aligned}
J^{(6)} &= \frac{8(n-1)}{n-2} \sum_{i,k=1}^n (g^{ik} - \delta_{ik} + H_{ik}) \partial_i u_\varepsilon \partial_k w \\
&\quad + 2 \left[R_g - \sum_{i,k=1}^n \partial_i \partial_k H_{ik} \right] u_\varepsilon w \\
J^{(7)} &= R_g w^2 + \frac{4(n-1)}{n-2} \sum_{i,k=1}^n (g^{ik} - \delta_{ik}) \partial_i w \partial_k w.
\end{aligned}$$

It follows from the divergence theorem that

$$\begin{aligned}
\int_{B_\delta(0) \cap \mathbb{R}_+^n} J^{(1)} &= \int_{B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \partial_i \left[\partial_i u_\varepsilon w + \frac{(n-2)^2}{2} u_\varepsilon^{\frac{2n}{n-2}} V_i \right] \\
&= \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} \left[\partial_i u_\varepsilon w + \frac{(n-2)^2}{2} u_\varepsilon^{\frac{2n}{n-2}} V_i \right] \\
&\leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2}.
\end{aligned}$$

We next observe that

$$\begin{aligned}
J^{(2)} &= \sum_{i,k=1}^n \partial_i (u_\varepsilon^2 \partial_k h_{ik} - 2 u_\varepsilon \partial_k u_\varepsilon h_{ik}) \\
&= 2 \sum_{i,k=1}^n \left(u_\varepsilon \partial_i \partial_k u_\varepsilon - \frac{n}{n-2} \partial_i u_\varepsilon \partial_k u_\varepsilon \right) h_{ik} \\
&= \frac{2}{n} \left(u_\varepsilon \Delta u_\varepsilon - \frac{n}{n-2} |du_\varepsilon|^2 \right) \operatorname{tr} h \\
&\leq C \varepsilon^{n-2} (\varepsilon + |x|)^{2d+4-2n}.
\end{aligned}$$

Using the divergence theorem, we obtain

$$\begin{aligned}
&\int_{B_\delta(0) \cap \mathbb{R}_+^n} J^{(2)} \\
&\leq \sum_{i,k=1}^n \partial_i (u_\varepsilon^2 \partial_k h_{ik} - 2 u_\varepsilon \partial_k u_\varepsilon h_{ik}) + C \delta^{2d+4-n} \varepsilon^{n-2} \\
&\leq \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k=1}^n \frac{x_i}{|x|} (u_\varepsilon^2 \partial_k h_{ik} - 2 u_\varepsilon \partial_k u_\varepsilon h_{ik}) + C \delta^{2d+4-n} \varepsilon^{n-2}.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
\int_{B_\delta(0) \cap \mathbb{R}_+^n} J^{(3)} &= - \int_{B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n \partial_k (u_\varepsilon^2 H_{ik} \partial_l H_{il}) \\
&= - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k,l=1}^n \frac{x_k}{|x|} u_\varepsilon^2 H_{ik} \partial_l H_{il} \\
&\leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{n-2}.
\end{aligned}$$

Using Proposition 12, we obtain

$$\begin{aligned}
&\int_{B_\delta(0) \cap \mathbb{R}_+^n} J^{(4)} \\
&\leq -2\lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
&+ C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{n-2}.
\end{aligned}$$

It remains to estimate the terms $J^{(5)}$, $J^{(6)}$, and $J^{(7)}$. Using Proposition 13, we obtain the pointwise estimate

$$\begin{aligned}
J^{(5)} + J^{(6)} + J^{(7)} &\leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} (\varepsilon + |x|)^{2|\alpha|+4-2n} \\
&+ C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \varepsilon^{n-2} (\varepsilon + |x|)^{|\alpha|+d+3-2n} \\
&+ C \varepsilon^{n-2} (\varepsilon + |x|)^{2d+4-2n}
\end{aligned}$$

for $x \in B_\delta(0) \cap \mathbb{R}_+^n$. Using Young's inequality, we deduce that

$$\begin{aligned}
J^{(5)} + J^{(6)} + J^{(7)} &\leq \lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} (\varepsilon + |x|)^{2|\alpha|+2-2n} \\
&+ C \varepsilon^{n-2} (\varepsilon + |x|)^{2d+4-2n}
\end{aligned}$$

for $x \in B_\delta(0) \cap \mathbb{R}_+^n$. Integration over $B_\delta(0) \cap \mathbb{R}_+^n$ yields

$$\begin{aligned}
&\int_{B_\delta(0) \cap \mathbb{R}_+^n} (J^{(5)} + J^{(6)} + J^{(7)}) \\
&\leq \lambda \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
&+ C \delta^{2d+4-n} \varepsilon^{n-2}.
\end{aligned}$$

Putting these facts together, the assertion follows.

Proposition 15. *If δ_0 is sufficiently small, then we have*

$$\begin{aligned}
& \int_{B_\delta(0) \cap \mathbb{R}_+^n} \left(u_\varepsilon^2 + \frac{n+2}{n-2} w^2 \right)^{\frac{n}{n-2}} \\
& \leq \int_{B_\delta(0) \cap \mathbb{R}_+^n} (u_\varepsilon + w)^{\frac{2n}{n-2}} \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^n \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \delta^{|\alpha|-n} \varepsilon^n
\end{aligned}$$

for all $0 < 2\varepsilon \leq \delta \leq \delta_0$.

Proof. The proof is analogous to the proof of Proposition 14 in [4]. We omit the details.

Proposition 16. *If δ_0 is sufficiently small, then we have*

$$\begin{aligned}
& \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} |du_\varepsilon|^2 + \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
& \leq Y(S_+^n, \partial S_+^n) \left(\int_{B_\delta(0) \cap \mathbb{R}_+^n} (u_\varepsilon + w)^{\frac{2n}{n-2}} \right)^{\frac{n-2}{n}} \\
& + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 \varepsilon^n \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| \delta^{|\alpha|-n} \varepsilon^n
\end{aligned}$$

for all $0 < 2\varepsilon \leq \delta \leq \delta_0$.

Proof. The proof is similar to the proof of Proposition 15 in [4]. We first observe that

$$4n(n-1) \left(\int_{\mathbb{R}_+^n} u_\varepsilon^{\frac{2n}{n-2}} \right)^{\frac{2}{n}} = Y(S_+^n, \partial S_+^n).$$

Using Hölder's inequality, we obtain

$$\begin{aligned}
& \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} |du_\varepsilon|^2 - \int_{\partial B_\delta(0)} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon \\
& + \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
& = - \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \Delta u_\varepsilon u_\varepsilon + \int_{B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} n(n+2) u_\varepsilon^{\frac{4}{n-2}} w^2 \\
& = \int_{B_\delta(0) \cap \mathbb{R}_+^n} 4n(n-1) u_\varepsilon^{\frac{4}{n-2}} \left(u_\varepsilon^2 + \frac{n+2}{n-2} w^2 \right) \\
& \leq Y(S_+^n, \partial S_+^n) \left(\int_{B_\delta(0) \cap \mathbb{R}_+^n} \left(u_\varepsilon^2 + \frac{n+2}{n-2} w^2 \right)^{\frac{n-2}{n}} \right)^{\frac{n}{n-2}}.
\end{aligned}$$

Hence, the assertion follows from Proposition 15.

4. PROOF OF THE MAIN RESULT

In this section, we construct a smooth function $v_{(\varepsilon, \delta)} : M \rightarrow \mathbb{R}$ with Yamabe energy less than $Y(S_+^n, \partial S_+^n)$. The existence of such a function is trivial when $Y(M, \partial M, g) \leq 0$. Hence, it suffices to consider the case $Y(M, \partial M, g) > 0$. As in the previous section, we fix a boundary point $p \in \partial M$. Moreover, we denote by $G : M \setminus \{p\} \rightarrow \mathbb{R}$ the Greens function for the conformal Laplacian with Neumann boundary condition with pole at p . In other words, G satisfies

$$\frac{4(n-1)}{n-2} \Delta_g G - R_g G = 0$$

in $M \setminus \{p\}$ and $\partial_\nu G = 0$ along $\partial M \setminus \{p\}$. We assume that $G_p(x)$ is normalized so that $\lim_{x \rightarrow 0} |x|^{n-2} G(x) = 1$. With this normalization, we have

$$(7) \quad |G(x) - |x|^{2-n}| \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}| |x|^{|\alpha|+2-n} + C |x|^{d+3-n}.$$

Moreover, we consider the flux integral

$$\begin{aligned}
\mathcal{I}(p, \delta) &= \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} (|x|^{2-n} \partial_i G - G \partial_i |x|^{2-n}) \\
&\quad - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} |x|^{1-2n} \sum_{i,k=1}^n x_i (|x|^2 \partial_k h_{ik} - 2n x_k h_{ik})
\end{aligned}$$

where $\delta > 0$ is sufficiently small.

We next define a function $v_{(\varepsilon, \delta)} : M \rightarrow \mathbb{R}$ by

$$(8) \quad v_{(\varepsilon, \delta)} = \chi_\delta (u_\varepsilon + w) + (1 - \chi_\delta) \varepsilon^{\frac{n-2}{2}} G,$$

where χ_δ is the cut-off function defined above. Our main result is an upper bound for the Yamabe energy of $v_{(\varepsilon, \delta)}$:

Proposition 17. *If δ_0 is sufficiently small, then we have*

$$\begin{aligned}
& \int_M \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& \leq Y(S_+^n, \partial S_+^n) \left(\int_M v_{(\varepsilon, \delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}} - \varepsilon^{n-2} \mathcal{I}(p, \delta) \\
& \quad - \frac{\lambda}{2} \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& \quad + C \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2} + C \delta^{-n} \varepsilon^n
\end{aligned}$$

for all $0 < 2\varepsilon \leq \delta \leq \delta_0$.

Proof. For abbreviation, we denote by Ω_δ the set of all points in M such that $x_1^2 + \dots + x_n^2 < \delta^2$, where $(x_1, \dots, x_n)$ denote the Fermi coordinates around p . (In other words, Ω_δ is a coordinate ball, not a geodesic ball.) Using the divergence theorem, we obtain

$$\begin{aligned}
& \int_{M \setminus \Omega_\delta} \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& = - \int_{M \setminus \Omega_\delta} \left(\frac{4(n-1)}{n-2} \Delta_g v_{(\varepsilon, \delta)} - R_g v_{(\varepsilon, \delta)} \right) (v_{(\varepsilon, \delta)} - \varepsilon^{\frac{n-2}{2}} G) d\text{vol}_g \\
& \quad - \int_{\partial \Omega_\delta} \frac{4(n-1)}{n-2} \partial_\nu v_{(\varepsilon, \delta)} v_{(\varepsilon, \delta)} d\sigma_g \\
& \quad - \int_{\partial \Omega_\delta} \frac{4(n-1)}{n-2} \varepsilon^{\frac{n-2}{2}} (v_{(\varepsilon, \delta)} \partial_\nu G - G \partial_\nu v_{(\varepsilon, \delta)}) d\sigma_g,
\end{aligned}$$

where ν denotes the outward-pointing unit normal to $\partial \Omega_\delta$. Note that

$$v_{(\varepsilon, \delta)} - \varepsilon^{\frac{n-2}{2}} G = \chi_\delta (u_\varepsilon + w - \varepsilon^{\frac{n-2}{2}} G)$$

in $M \setminus \Omega_\delta$. In particular, we have $v_{(\varepsilon, \delta)} - \varepsilon^{\frac{n-2}{2}} G = 0$ in $M \setminus \Omega_{2\delta}$. Using (7), we obtain

$$\begin{aligned}
& \sup_{M \setminus \Omega_\delta} |v_{(\varepsilon, \delta)} - \varepsilon^{\frac{n-2}{2}} G| + \delta^2 \sup_{M \setminus \Omega_\delta} \left| \frac{4(n-1)}{n-2} \Delta_g v_{(\varepsilon, \delta)} - R_g v_{(\varepsilon, \delta)} \right| \\
& \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}| \delta^{|\alpha|+2-n} \varepsilon^{\frac{n-2}{2}} + C \delta^{d+3-n} \varepsilon^{\frac{n-2}{2}} + C \delta^{-n} \varepsilon^{\frac{n+2}{2}},
\end{aligned}$$

hence

$$\begin{aligned}
& - \int_{M \setminus \Omega_\delta} \left(\frac{4(n-1)}{n-2} \Delta_g v_{(\varepsilon, \delta)} - R_g v_{(\varepsilon, \delta)} \right) (v_{(\varepsilon, \delta)} - \varepsilon^{\frac{n-2}{2}} G) d\text{vol}_g \\
& \leq C \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2} + C \delta^{-n-2} \varepsilon^{n+2}.
\end{aligned}$$

We next observe that

$$\begin{aligned}
& - \int_{\partial \Omega_\delta} \partial_\nu u_\varepsilon u_\varepsilon d\sigma_g \\
& \leq - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} u_\varepsilon \partial_k u_\varepsilon h_{ik} \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2},
\end{aligned}$$

hence

$$\begin{aligned}
& - \int_{\partial \Omega_\delta} \partial_\nu v_{(\varepsilon, \delta)} v_{(\varepsilon, \delta)} d\sigma_g \\
& \leq - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} u_\varepsilon \partial_k u_\varepsilon h_{ik} \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i, k=1}^n |h_{ik, \alpha}|^2 \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2}.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
& - \int_{\partial \Omega_\delta} (v_{(\varepsilon, \delta)} \partial_\nu G - G \partial_\nu v_{(\varepsilon, \delta)}) d\sigma_g \\
& \leq - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i=1}^n \frac{x_i}{|x|} (u_\varepsilon \partial_i G - G \partial_i u_\varepsilon) \\
& + C \sum_{2 \leq |\alpha| \leq d} |h_{ik, \alpha}|^2 \delta^{2|\alpha|+2-n} \varepsilon^{\frac{n-2}{2}} + C \delta^{2d+4-n} \varepsilon^{\frac{n-2}{2}}.
\end{aligned}$$

Putting these facts together, we obtain

$$\begin{aligned}
& \int_{M \setminus \Omega_\delta} \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& \leq - \int_{\partial\Omega_\delta} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} u_\varepsilon \partial_k u_\varepsilon h_{ik} \\
& \quad - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \varepsilon^{\frac{n-2}{2}} \sum_{i=1}^n \frac{x_i}{|x|} (u_\varepsilon \partial_i G - G \partial_i u_\varepsilon) \\
& \quad + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2} + C \delta^{-n-2} \varepsilon^{n+2}.
\end{aligned}$$

On the other hand, it follows from Proposition 14 and Proposition 16 that

$$\begin{aligned}
& \int_{\Omega_\delta} \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& \leq Y(S_+^n, \partial S_+^n) \left(\int_{\Omega_\delta} v_{(\varepsilon, \delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}} + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{x_i}{|x|} \partial_i u_\varepsilon u_\varepsilon \\
& \quad + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k=1}^n \frac{x_i}{|x|} (u_\varepsilon^2 \partial_k h_{ik} - 2 u_\varepsilon \partial_k u_\varepsilon h_{ik}) \\
& \quad - \frac{\lambda}{2} \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& \quad + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2}.
\end{aligned}$$

If we add the last two inequalities, we obtain

$$\begin{aligned}
& \int_M \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& \leq Y(S_+^n, \partial S_+^n) \left(\int_{\Omega_\delta} v_{(\varepsilon, \delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}} \\
& + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \sum_{i,k=1}^n \frac{x_i}{|x|} \left(u_\varepsilon^2 \partial_k h_{ik} + \frac{2n}{n-2} u_\varepsilon \partial_k u_\varepsilon h_{ik} \right) \\
& - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} \varepsilon^{\frac{n-2}{2}} \sum_{i=1}^n \frac{x_i}{|x|} (u_\varepsilon \partial_i G - G \partial_i u_\varepsilon) \\
& - \frac{\lambda}{2} \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}|^2 \varepsilon^{n-2} \int_{B_\delta(0) \cap \mathbb{R}_+^n} (\varepsilon + |x|)^{2|\alpha|+2-2n} dx \\
& + C \sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}| \delta^{|\alpha|+2-n} \varepsilon^{n-2} + C \delta^{2d+4-n} \varepsilon^{n-2} + C \delta^{-n-2} \varepsilon^{n+2}.
\end{aligned}$$

From this, the assertion follows easily.

Theorem 18. *Assume that $p \notin \mathcal{Z}$. Then $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$.*

Proof. Since $p \notin \mathcal{Z}$, we have $\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik, \alpha}|^2 > 0$. Using Proposition 17, we obtain

$$\begin{aligned}
& \int_M \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\
& < Y(S_+^n, \partial S_+^n) \left(\int_M v_{(\varepsilon, \delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}}
\end{aligned}$$

if $\varepsilon > 0$ is sufficiently small. From this, the assertion follows.

In the remainder of this section, we study the case $p \in \mathcal{Z}$. In this case, we consider the manifold $(M \setminus \{p\}, G^{\frac{4}{n-2}} g)$. This manifold is scalar flat and its boundary is totally geodesic. After doubling this manifold, we obtain an asymptotically flat manifold with zero scalar curvature.

Proposition 19. *Assume that $p \in \mathcal{Z}$. Then the following statements hold:*

- (i) *The limit $\lim_{\delta \rightarrow 0} \mathcal{I}(p, \delta)$ exists.*
- (ii) *The doubling of $(M \setminus \{p\}, G^{\frac{4}{n-2}} g)$ has a well-defined mass which equals $\lim_{\delta \rightarrow 0} \mathcal{I}(p, \delta)$ up to a positive factor.*

Proof. For abbreviation, let $\bar{g} = G^{\frac{4}{n-2}} g$. We consider the inverted coordinates $y = \frac{x}{|x|^2}$, where $(x_1, \dots, x_n)$ are conformal Fermi coordinates

around p . In these coordinates, the metric $\bar{g}$ is given by

$$\begin{aligned} \bar{g}\left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l}\right) &= \left[1 + \Phi\left(\frac{y}{|y|^2}\right)\right]^{\frac{4}{n-2}} \\ &\quad \cdot |y|^{-4} \sum_{i,k=1}^n (|y|^2 \delta_{ij} - 2y_i y_j) (|y|^2 \delta_{kl} - 2y_k y_l) g_{ik}\left(\frac{y}{|y|^2}\right), \end{aligned}$$

where $\Phi(x) = |x|^{n-2} G(x) - 1$. Using the relations $g_{ik}(x) = \delta_{ik} + h_{ik}(x) + O(|x|^{2d+2})$ and $\Phi(x) = O(|x|^{d+1})$, we obtain

$$\begin{aligned} \bar{g}\left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l}\right) &= \left[1 + \frac{4}{n-2} \Phi\left(\frac{y}{|y|^2}\right)\right] \delta_{jl} \\ &\quad + |y|^{-4} \sum_{i,k=1}^n (|y|^2 \delta_{ij} - 2y_i y_j) (|y|^2 \delta_{kl} - 2y_k y_l) h_{ik}\left(\frac{y}{|y|^2}\right) \\ &\quad + O(|y|^{-2d-2}). \end{aligned}$$

In particular, we have $\bar{g}\left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l}\right) = \delta_{jl} + O(|y|^{-d-1})$. Hence, the doubling of $(M \setminus \{p\}, \bar{g})$ is asymptotically flat in the sense of Bartnik [2], and has a well-defined ADM mass.

Since $\text{tr } h = O(|x|^{2d+2})$, it follows that

$$\sum_{j,l=1}^n y_j \frac{\partial}{\partial y_j} \bar{g}\left(\frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_l}\right) = -\frac{4n}{n-2} \sum_{j=1}^n \frac{y_j}{|y|^2} (\partial_j \Phi)\left(\frac{y}{|y|^2}\right) + O(|y|^{-2d-2}).$$

Moreover, we have

$$\begin{aligned} \sum_{j,l=1}^n y_j \frac{\partial}{\partial y_l} \bar{g}\left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l}\right) &= -\frac{4}{n-2} \sum_{i=1}^n \frac{y_i}{|y|^2} (\partial_i \Phi)\left(\frac{y}{|y|^2}\right) \\ &\quad - \sum_{i,k=1}^n \frac{y_i}{|y|^2} (\partial_k h_{ik})\left(\frac{y}{|y|^2}\right) \\ &\quad + 2n \sum_{i,k=1}^n \frac{y_i y_k}{|y|^2} h_{ik}\left(\frac{y}{|y|^2}\right) + O(|y|^{-2d-2}), \end{aligned}$$

where $\partial_i \Phi(x) = \frac{\partial}{\partial x_i} \Phi(x)$. Putting these facts together, we obtain

$$\begin{aligned} & \sum_{j,l=1}^n y_j \frac{\partial}{\partial y_l} \bar{g} \left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l} \right) - \sum_{j,l=1}^n y_j \frac{\partial}{\partial y_j} \bar{g} \left(\frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_l} \right) \\ &= \frac{4(n-1)}{n-2} \sum_{i=1}^n \frac{y_i}{|y|^2} (\partial_i \Phi) \left(\frac{y}{|y|^2} \right) \\ & - \sum_{i,k=1}^n \frac{y_i}{|y|^2} (\partial_k h_{ik}) \left(\frac{y}{|y|^2} \right) + 2n \sum_{i,k=1}^n \frac{y_i y_k}{|y|^2} h_{ik} \left(\frac{y}{|y|^2} \right) \\ & + O(|y|^{-2d-2}). \end{aligned}$$

This implies

$$\begin{aligned} & \int_{\partial B_{\frac{1}{\delta}}(0) \cap \mathbb{R}_+^n} \sum_{j,l=1}^n \frac{y_j}{|y|} \frac{\partial}{\partial y_l} \bar{g} \left(\frac{\partial}{\partial y_j}, \frac{\partial}{\partial y_l} \right) - \int_{\partial B_{\frac{1}{\delta}}(0) \cap \mathbb{R}_+^n} \sum_{j,l=1}^n \frac{y_j}{|y|} \frac{\partial}{\partial y_j} \bar{g} \left(\frac{\partial}{\partial y_l}, \frac{\partial}{\partial y_l} \right) \\ &= \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} \frac{4(n-1)}{n-2} |x|^{3-2n} \sum_{i=1}^n x_i \partial_i \Phi(x) \\ & - \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} |x|^{3-2n} \sum_{i,k=1}^n x_i (\partial_k h_{ik})(x) + \int_{\partial B_\delta(0) \cap \mathbb{R}_+^n} 2n |x|^{1-2n} \sum_{i,k=1}^n x_i x_k h_{ik}(x) \\ & + O(\delta^{2d+4-n}) \\ & = \mathcal{I}(p, \delta) + O(\delta^{2d+n-4}). \end{aligned}$$

As $\delta \rightarrow 0$, the left hand side converges to a positive multiple of the ADM mass. From this the assertion follows.

Theorem 20. *Assume that $p \in \mathcal{Z}$. If $\lim_{\delta \rightarrow 0} \mathcal{I}(p, \delta)$ is positive, then $Y(M, \partial M, g) < Y(S_+^n, \partial S_+^n)$.*

Proof. Since $p \in \mathcal{Z}$, we have $\sum_{2 \leq |\alpha| \leq d} \sum_{i,k=1}^n |h_{ik,\alpha}|^2 = 0$. By Proposition 17, we can find positive real numbers δ_0 and C such that

$$\begin{aligned} & \int_M \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon,\delta)}|_g^2 + R_g v_{(\varepsilon,\delta)}^2 \right) d\text{vol}_g \\ & \leq Y(S_+^n, \partial S_+^n) \left(\int_M v_{(\varepsilon,\delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}} - \varepsilon^{n-2} \mathcal{I}(p, \delta) \\ & + C \delta^{2d+4-n} \varepsilon^{n-2} + C \delta^{-n} \varepsilon^n \end{aligned}$$

whenever $0 < 2\varepsilon \leq \delta \leq \delta_0$. Since $\lim_{\delta \rightarrow 0} \mathcal{I}(p, \delta)$ is positive, we can find a real number $\delta \in (0, \delta_0]$ such that $\mathcal{I}(p, \delta) > C \delta^{2d+4-n}$. In the next step, we choose $\varepsilon \in (0, \frac{\delta}{2}]$ small enough so that $\mathcal{I}(p, \delta) > C \delta^{2d+4-n} + C \delta^{-n} \varepsilon^2$. For

this choice of ε and δ , we have

$$\begin{aligned} & \int_M \left(\frac{4(n-1)}{n-2} |dv_{(\varepsilon, \delta)}|_g^2 + R_g v_{(\varepsilon, \delta)}^2 \right) d\text{vol}_g \\ & < Y(S_+^n, \partial S_+^n) \left(\int_M v_{(\varepsilon, \delta)}^{\frac{2n}{n-2}} d\text{vol}_g \right)^{\frac{n-2}{n}}. \end{aligned}$$

This completes the proof.

APPENDIX A. AN ELLIPTIC SYSTEM ON $\mathbb{R}_+^n$

In this section, we describe the construction of the vector field V . In the following, we consider the hemisphere S_+^n , equipped with the round metric of constant sectional curvature 4. We denote by $\mathcal{X}$ the space of all vector fields V on S_+^n such that V is of class H^1 and $\langle V, \nu \rangle = 0$ along ∂S_+^n . Moreover, we denote by $\mathcal{Y}$ the space of all trace-free symmetric two-tensors on S_+^n of class L^2 . We next define a linear operator $\mathcal{D} : \mathcal{X} \rightarrow \mathcal{Y}$ by

$$\mathcal{D}V = \widehat{\mathcal{L}}_V g = \mathcal{L}_V g - \frac{2}{n} (\text{div}_g V) g.$$

In other words, $\mathcal{D}$ is the conformal Killing operator.

Lemma 21. *We have*

$$\|\nabla V\|_{L^2(S_+^n)}^2 \leq \|\mathcal{D}V\|_{L^2(S_+^n)}^2 + 4(n-1) \|V\|_{L^2(S_+^n)}^2$$

for all $V \in \mathcal{X}$.

Proof. Without loss of generality, we may assume that V is smooth. By definition of $\mathcal{D}$, we have

$$\|\mathcal{D}V\|_{L^2(S_+^n)}^2 = \int_{S_+^n} \left[\nabla_i V^k \nabla^i V_k + \nabla_i V^k \nabla_k V^i - \frac{2}{n} (\text{div}_g V)^2 \right] d\text{vol}_g.$$

Integration by parts yields

$$\begin{aligned} \int_{S_+^n} \nabla_i V^k \nabla_k V^i d\text{vol}_g &= - \int_{S_+^n} V^k \nabla_i \nabla_k V^i d\text{vol}_g \\ &= - \int_{S_+^n} V^k \nabla_k \nabla_i V^i d\text{vol}_g - \int_{S_+^n} \text{Ric}_{ik} V^i V^k d\text{vol}_g \\ &= \int_{S_+^n} (\text{div}_g V)^2 d\text{vol}_g - 4(n-1) \int_{S_+^n} |V|^2 d\text{vol}_g. \end{aligned}$$

Putting these facts together, we obtain

$$\|\mathcal{D}V\|_{L^2(S_+^n)}^2 + 4(n-1) \|V\|_{L^2(S_+^n)}^2 = \|\nabla V\|_{L^2(S_+^n)}^2 + \frac{n-2}{n} \|\text{div}_g V\|_{L^2(S_+^n)}^2.$$

From this, the assertion follows.

It follows from Lemma 21 and Rellich's theorem that $\ker \mathcal{D}$ is finite-dimensional. We now consider the subspace

$$\mathcal{X}_0 = \{V \in \mathcal{X} : \langle V, W \rangle_{L^2(S_+^n)} = 0 \text{ for all } W \in \ker \mathcal{D}\}.$$

Lemma 22. *We have*

$$\|V\|_{L^2(S_+^n)}^2 + \|\nabla V\|_{L^2(S_+^n)}^2 \leq K \|\mathcal{D}V\|_{L^2(S_+^n)}^2$$

for all $V \in \mathcal{X}_0$. Here, K is a positive constant that depends only on n .

Proof. Suppose that the assertion is false. Then we can find a sequence of vector fields $V^{(\nu)} \in \mathcal{X}_0$ such that

$$(9) \quad \|V^{(\nu)}\|_{L^2(S_+^n)}^2 + \|\nabla V^{(\nu)}\|_{L^2(S_+^n)}^2 = 1$$

for all ν and $\|\mathcal{D}V^{(\nu)}\|_{L^2(S_+^n)} \rightarrow 0$ as $\nu \rightarrow \infty$. After passing to a subsequence, we may assume that the sequence $V^{(\nu)}$ converges weakly to a vector field $W \in \mathcal{X}_0$. Then $\mathcal{D}W = 0$. Since $W \in \mathcal{X}_0$, we conclude that $W = 0$. This implies $\|V^{(\nu)}\|_{L^2(S_+^n)} \rightarrow 0$ as $\nu \rightarrow \infty$. Using Lemma 21, we obtain $\|\nabla V^{(\nu)}\|_{L^2(S_+^n)} \rightarrow 0$ as $\nu \rightarrow \infty$. This contradicts (9).

Proposition 23. *Given any $h \in \mathcal{Y}$, there exists a unique vector field $V \in \mathcal{X}_0$ such that $\langle h - \mathcal{D}V, \mathcal{D}W \rangle_{L^2(S_+^n)} = 0$ for all $W \in \mathcal{X}$. The vector field V satisfies the estimate*

$$(10) \quad \|V\|_{L^2(S_+^n)}^2 + \|\nabla V\|_{L^2(S_+^n)}^2 \leq K \|h\|_{L^2(S_+^n)}^2.$$

Proof. It follows from Lemma 22 that the operator $\mathcal{D} : \mathcal{X}_0 \rightarrow \mathcal{Y}$ has closed range. Hence, we can find a vector field $V \in \mathcal{X}_0$ such that $\|h - \mathcal{D}V\|_{L^2(S_+^n)}^2$ is minimal. This vector field satisfies $\langle h - \mathcal{D}V, \mathcal{D}W \rangle_{L^2(S_+^n)} = 0$ for all $W \in \mathcal{X}_0$. This proves the existence statement.

We next assume that $V \in \mathcal{X}_0$ is a vector field satisfying $\langle h - \mathcal{D}V, \mathcal{D}W \rangle_{L^2(S_+^n)} = 0$ for all $W \in \mathcal{X}_0$. This implies $\langle h - \mathcal{D}V, \mathcal{D}V \rangle = 0$, hence $\|\mathcal{D}V\|_{L^2(S_+^n)}^2 \leq \|h\|_{L^2(S_+^n)}^2$. Thus, we conclude that

$$\|V\|_{L^2(S_+^n)}^2 + \|\nabla V\|_{L^2(S_+^n)}^2 \leq K \|\mathcal{D}V\|_{L^2(S_+^n)}^2 \leq K \|h\|_{L^2(S_+^n)}^2$$

by Lemma 22. In particular, if $h = 0$, then $V = 0$. From this, the uniqueness statement follows.

In the next step, we consider the stereographic projection from S_+^n to $\mathbb{R}_+^n \cup \{\infty\}$. The metric g can be written in the form $g_{ik} = u^{\frac{4}{n-2}} \delta_{ik}$, where

$$u(x) = \left(\frac{1}{1 + |x|^2} \right)^{\frac{n-2}{2}}.$$

Theorem 24. *Let h be a trace-free symmetric two-tensor on $\mathbb{R}_+^n$. We assume that h is smooth and has compact support. Then there exists a smooth vector field V on $\mathbb{R}_+^n$ with the following properties:*

- *At each point $x \in \mathbb{R}_+^n$, we have*

$$\sum_{k=1}^n \partial_k \left[u^{\frac{2n}{n-2}} (h_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}) \right] = 0$$

for all $i \in \{1, \dots, n\}$.

- *At each point $x \in \partial\mathbb{R}_+^n$, we have $V_n(x) = \partial_n V_i(x) - h_{in}(x) = 0$ for all $i \in \{1, \dots, n-1\}$.*

Moreover, the vector field V satisfies

$$(11) \quad \int_{\mathbb{R}_+^n} u(x)^{\frac{2(n+2)}{n-2}} |V(x)|^2 dx \leq K \int_{\mathbb{R}_+^n} u(x)^{\frac{2n}{n-2}} |h(x)|^2 dx.$$

Proof. By Proposition 23, there exists a smooth vector field $V \in \mathcal{X}_0$ such that

$$\int_{\mathbb{R}_+^n} \langle u^{\frac{4}{n-2}} h - \mathcal{D}V, \mathcal{D}W \rangle_g d\operatorname{vol}_g = 0$$

for all vector fields $W \in \mathcal{X}$. This implies

$$(12) \quad \int_{\mathbb{R}_+^n} u^{\frac{2n}{n-2}} \sum_{i,k=1}^n (h_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}) \partial_k W_i dx = 0$$

for all $W \in \mathcal{X}$. Since $V \in \mathcal{X}_0$, we have $V_n(x) = 0$ for $x \in \partial\mathbb{R}_+^n$.

By assumption, h is smooth. Using general regularity results for elliptic systems (cf. [10], [13]), we conclude that V is smooth. Using (12), we obtain

$$\sum_{k=1}^n \partial_k \left[u^{\frac{2n}{n-2}} (h_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}) \right] = 0$$

for all points $x \in \mathbb{R}_+^n$ and all $i \in \{1, \dots, n\}$. Moreover, at each point $x \in \partial\mathbb{R}_+^n$, we have $\partial_n V_i(x) - h_{in}(x) = 0$ for $i \in \{1, \dots, n-1\}$. Finally, the estimate (11) follows immediately from (10).

Proposition 25. *Fix a real number σ such that $1 < \sigma < n-2$. Let h be a trace-free symmetric two-tensor on $\mathbb{R}_+^n$ which is smooth and has compact support. Moreover, let V be the vector field constructed in Theorem 24. Finally, let us assume that*

$$\sup_{r \geq 1} r^{-2\sigma-n-2} \int_{(B_{2r}(0) \setminus B_r(0)) \cap \mathbb{R}_+^n} |V(x)|^2 dx < \infty.$$

Then there exists a constant C , depending only on n and σ , such that

$$\begin{aligned}
 (13) \quad & \sup_{r \geq 1} r^{-2\sigma-n-2} \int_{(B_{2r}(0) \setminus B_r(0)) \cap \mathbb{R}_+^n} |V(x)|^2 dx \\
 & \leq C \int_{\mathbb{R}_+^n} (1 + |x|^2)^{-n-2} |V(x)|^2 dx \\
 & + C \sup_{r \geq 1} r^{-2\sigma-n} \int_{(B_{2r}(0) \setminus B_r(0)) \cap \mathbb{R}_+^n} |h(x)|^2 dx.
 \end{aligned}$$

Proof. We extend V and h to $\mathbb{R}^n$ by reflection. More precisely, we define a vector field $\tilde{V}$ on $\mathbb{R}^n$ by

$$\begin{aligned}
 \tilde{V}_i(x_1, \dots, x_{n-1}, x_n) &= \tilde{V}_i(x_1, \dots, x_{n-1}, -x_n) = V_i(x_1, \dots, x_{n-1}, x_n) \\
 \tilde{V}_n(x_1, \dots, x_{n-1}, x_n) &= -\tilde{V}_n(x_1, \dots, x_{n-1}, -x_n) = V_n(x_1, \dots, x_{n-1}, x_n)
 \end{aligned}$$

for all $x \in \mathbb{R}_+^n$ and all $i \in \{1, \dots, n-1\}$. Similarly, we define a trace-free symmetric two-tensor $\tilde{h}$ on $\mathbb{R}^n$ by

$$\begin{aligned}
 \tilde{h}_{ik}(x_1, \dots, x_{n-1}, x_n) &= \tilde{h}_{ik}(x_1, \dots, x_{n-1}, -x_n) = h_{ik}(x_1, \dots, x_{n-1}, x_n) \\
 \tilde{h}_{in}(x_1, \dots, x_{n-1}, x_n) &= -\tilde{h}_{in}(x_1, \dots, x_{n-1}, -x_n) = h_{in}(x_1, \dots, x_{n-1}, x_n) \\
 \tilde{h}_{nk}(x_1, \dots, x_{n-1}, x_n) &= -\tilde{h}_{nk}(x_1, \dots, x_{n-1}, -x_n) = h_{nk}(x_1, \dots, x_{n-1}, x_n) \\
 \tilde{h}_{nn}(x_1, \dots, x_{n-1}, x_n) &= \tilde{h}_{nn}(x_1, \dots, x_{n-1}, -x_n) = h_{nn}(x_1, \dots, x_{n-1}, x_n)
 \end{aligned}$$

for all $x \in \mathbb{R}_+^n$ and all $i, k \in \{1, \dots, n-1\}$.

Since $V \in \mathcal{X}$, we have $V_n(x) = 0$ for all $x \in \partial\mathbb{R}_+^n$. Consequently, the vector field $\tilde{V}$ is a vector field on S^n of class H^1 . We claim that

$$(14) \quad \int_{\mathbb{R}^n} u^{\frac{2n}{n-2}} \sum_{i,k=1}^n (\tilde{h}_{ik} - \partial_i \tilde{V}_k - \partial_k \tilde{V}_i + \frac{2}{n} \operatorname{div} \tilde{V} \delta_{ik}) \partial_k \tilde{W}_i dx = 0$$

for all vector fields $\tilde{W}$ on S^n of class H^1 . In order to prove (14), we fix a vector field $\tilde{W}$ of class H^1 . We then define a vector field W on S_+^n by

$$\begin{aligned}
 W_i(x_1, \dots, x_{n-1}, x_n) &= \tilde{W}_i(x_1, \dots, x_{n-1}, x_n) + \tilde{W}_i(x_1, \dots, x_{n-1}, -x_n) \\
 W_n(x_1, \dots, x_{n-1}, x_n) &= \tilde{W}_n(x_1, \dots, x_{n-1}, x_n) - \tilde{W}_n(x_1, \dots, x_{n-1}, -x_n)
 \end{aligned}$$

for all $x \in \mathbb{R}_+^n$ and all $i \in \{1, \dots, n-1\}$. Clearly, $W \in \mathcal{X}$. Therefore, we have

$$\int_{\mathbb{R}_+^n} u^{\frac{2n}{n-2}} \sum_{i,k=1}^n (h_{ik} - \partial_i V_k - \partial_k V_i + \frac{2}{n} \operatorname{div} V \delta_{ik}) \partial_k W_i dx = 0$$

by definition of V . From this, the identity (14) follows easily.

We now complete the proof of Proposition 25. Using Proposition 23 in [4], we obtain

$$\begin{aligned} & \sup_{r \geq 1} r^{-2\sigma-n-2} \int_{B_{2r}(0) \setminus B_r(0)} |\tilde{V}(x)|^2 dx \\ & \leq C \int_{\mathbb{R}_+^n} (1 + |x|^2)^{-n-2} |\tilde{V}(x)|^2 dx \\ & \quad + C \sup_{r \geq 1} r^{-2\sigma-n} \int_{B_{2r}(0) \setminus B_r(0)} |\tilde{h}(x)|^2 dx. \end{aligned}$$

Here, C is a positive constant that depends only on σ and n . (In [4], this result was stated in the special case that $\tilde{V}$ and $\tilde{h}$ are smooth, but the proof only requires that $\tilde{h}$ belongs to L^2 and $\tilde{V}$ is of class H^1 .) From this the assertion follows.

Corollary 26. *Consider a trace-free symmetric two-tensor of the form*

$$h_{ik}(x) = \chi(|x|/\rho) \sum_{2 \leq |\alpha| \leq d} h_{ik,\alpha} x^\alpha,$$

where $d = [\frac{n-2}{2}]$, $\rho \geq 1$, and $\chi : \mathbb{R} \rightarrow \mathbb{R}$ is a fixed cutoff function satisfying $\chi(t) = 0$ for $t \geq 2$. Let V be the vector field constructed in Theorem 24. Then, for every multi-index β , we have

$$(15) \quad |\partial^\beta V(x)|^2 \leq C \sum_{2 \leq |\alpha| \leq d} |h_{ik,\alpha}|^2 (1 + |x|^2)^{|\alpha|+1-|\beta|}$$

for all $x \in \mathbb{R}_+^n$. Here, C is positive constant which depends on n and $|\beta|$, but not on ρ .

Proof. Without loss of generality, we may assume that

$$h_{ik}(x) = \chi(|x|/\rho) \sum_{|\alpha|=d'} h_{ik,\alpha} x^\alpha,$$

where $2 \leq d' \leq d$. Since $d' < \frac{n}{2}$, we have

$$\int_{\mathbb{R}_+^n} (1 + |x|^2)^{-n} |h(x)|^2 dx \leq C \sum_{|\alpha|=d'} |h_{ik,\alpha}|^2$$

for some uniform constant C . Using (11), we obtain

$$\int_{\mathbb{R}_+^n} (1 + |x|^2)^{-n-2} |V(x)|^2 dx \leq C \sum_{|\alpha|=d'} |h_{ik,\alpha}|^2.$$

We now apply Proposition 25 with $\sigma = d'$. This yields

$$\sup_{r \geq 1} r^{-2d'-n-2} \int_{\{r \leq |x| \leq 2r\}} |V(x)|^2 dx \leq C \sum_{|\alpha|=d'} |h_{ik,\alpha}|^2.$$

Using elliptic estimates, we conclude that

$$|\partial^\beta V(x)|^2 \leq C \sum_{|\alpha|=d'} |h_{ik,\alpha}|^2 (1 + |x|^2)^{d'+1-|\beta|}$$

for every multi-index β .

REFERENCES

- [1] T. Aubin, *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*, J. Math. Pures Appl. 55, 269–296 (1976)
- [2] R. Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. 39, 661–693 (1986)
- [3] S. Brendle, *Convergence of the Yamabe flow for arbitrary initial energy*, J. Diff. Geom. 69, 217–278 (2005)
- [4] S. Brendle, *Convergence of the Yamabe flow in dimension 6 and higher*, Invent. Math. 170, 541–576 (2007)
- [5] S. Brendle, *Blow-up phenomena for the Yamabe equation*, J. Amer. Math. Soc. 21, 951–979 (2008)
- [6] S. Brendle and F. Marques, *Blow-up phenomena for the Yamabe equation II*, J. Diff. Geom. 81, 225–250 (2009)
- [7] S. Brendle, *On the conformal scalar curvature equation and related problems*, Surveys in Differential Geometry vol. XII, 1–19 (2008)
- [8] J. Escobar, *The Yamabe problem on manifolds with boundary*, J. Diff. Geom. 35, 21–84 (1992)
- [9] J. Escobar, *Conformal deformation of a Riemannian metric to a scalar flat metric with constant mean curvature at the boundary*, Ann. of Math. 136, 1–50 (1992)
- [10] L. Hörmander, *Linear partial differential operators*, Springer-Verlag, New York (1963)
- [11] F. Marques, *Existence results for the Yamabe problem on manifolds with boundary*, Indiana Univ. Math. J. 54, 1599–1620 (2005)
- [12] R.M. Schoen, *Conformal deformation of a Riemannian metric to constant scalar curvature*, J. Diff. Geom. 20, 479–495 (1984)
- [13] M. Taylor, *Partial differential equations III*, Springer-Verlag, New York (1997)
- [14] N. Trudinger, *Remarks concerning the conformal deformation of Riemannian structures on compact manifolds*, Annali Scuola Norm. Sup. Pisa 22, 265–274 (1968)

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305

SCHOOL OF MATHEMATICS, INSTITUTE FOR ADVANCED STUDY, PRINCETON, NJ 08540